# Waters™

Applikationsbericht

# Clopyralid and Triclopyr in River Water - LC/UV

Waters Corporation



This is an Application Brief and does not contain a detailed Experimental section.

## Abstract

This appliction brief highlights the analysis of clopyralid and triclopyr in river water using XTerra  $RP_{18}$  Columns.

## Introduction

Compounds used in this study are:

- 1. Clopyralid
- 2. Triclopyr

## Clopyralid

# Triclopyr

## Experimental

#### **HPLC Method**

Column:

XTerra RP<sub>18</sub> 4.6 x 100 mm, 3.5  $\mu$ m (p/n: 186000438)

Mobile phase A: 10 mM TFA, pH 2.1

Mobile phase B: ACN

Flow rate: 1.0 mL/min

Injection volume: 50  $\mu$ L


Detection: UV @ 290 nm

Instrument: Alliance 2695, 2996 PDA

## Gradient


Time (min)	Profile	
	%A	%B
0.0	80	20
20.0	20	80

#### Oasis® MAX Extraction Method Conditions for Oasis® MAX Cartridge, 6 cc, 500 mg Part Number 186000865



\* methyl #butyl ether diethyl ether can be used as an alternative to MTBE \*\* TFA - trifluoroacetic acid

## Results and Discussion



## Featured Products

Alliance HPLC <a href="https://www.waters.com/514248">https://www.waters.com/514248</a>

WA20738.026, June 2002

© 2021 Waters Corporation. All Rights Reserved.