

アプリケーションノート

Clopyralid and Triclopyr in River Water - LC/UV

Waters Corporation



This is an Application Brief and does not contain a detailed Experimental section.


Abstract

This application brief highlights the analysis of clopyralid and triclopyr in river water using XTerra RP₁₈ Columns.


Introduction

Compounds used in this study are:

1. Clopyralid
2. Triclopyr



Clopyralid



Triclopyr

Experimental

HPLC Method

Column:	XTerra RP ₁₈ 4.6 x 100 mm, 3.5 μm (p/n: 186000438)
Mobile phase A:	10 mM TFA, pH 2.1
Mobile phase B:	ACN
Flow rate:	1.0 mL/min

Injection volume:

50 μ L

Detection:

UV @ 290 nm


Instrument:

Alliance 2695, 2996 PDA

Gradient


Time (min)	Profile	
	%A	%B
0.0	80	20
20.0	20	80

Oasis® MAX Extraction Method
Conditions for Oasis® MAX Cartridge, 6 cc, 500 mg
Part Number 186000865



* methyl tbutyl ether
diethyl ether can be used as an alternative to MTBE
** TFA - trifluoroacetic acid

Results and Discussion



Featured Products

Alliance HPLC <<https://www.waters.com/514248>>

WA20738.026, June 2002