Waters™

응용 자료

Pseudoephedrine HCL and Triprolidine HCL

Waters Corporation



This is an Application Brief and does not contain a detailed Experimental section.

Abstract

This application brief highlights the analysis of Pseudoephedrine HCL and Triprolidine HCL using Symmetry Columns.

Introduction

The compounds analyzed in this study are:

- 1. Pseudoephedrine HCL
- 2. Triprolidine HCL

1. Pseudoephedrine HCL

$$\begin{array}{c|c} CH_{3} & & \\ \hline \\ C & C \\ \hline \\ H \\ \end{array} \begin{array}{c} CH_{2} - N \\ \hline \\ H \\ \end{array} \begin{array}{c} CH_{2} - N \\ \hline \\ H \\ \end{array} \begin{array}{c} CH_{2} - N \\ \hline \\ H \\ \end{array} \begin{array}{c} CH_{2} - N \\ \hline \\ H \\ \end{array}$$

2. Triprolidine HCL

Experimental

HPLC Method

Column:

Symmetry $C_8,\,3.9~x$ 150 mm, 5 μm

Guard column: Symmetry Guard Column 3.9 x 20 mm, 5 µm

Part numbers: Column - WAT046970, Guard - WAT054250

Mobile phase A: 50 mM potassium phosphate, pH 3.0

Mobile phase B: Acetonitrile

Flow rate: 1.0 mL/min

Injection volume: 5 μ L of 2.88 μ g/mL pseudoephedrine and 120 μ

g/mL triprolidine extracted tablet sample

Detection: UV @ 261 nm

Gradient Table


Time (min)	Profile	
	%A	%B
0	85	15
1	85	15
15	50	50

USP Tailing Factors

1. 1.53

2, 1,16

Results and Discussion



Featured Products

WA31763.138, June 2003

© 2021 Waters Corporation. All Rights Reserved.