Waters™

Note d'application

Separation of Nucleotide Phosphates on ACQUITY UPLC BEH Amide

Waters Corporation



This is an Application Brief and does not contain a detailed Experimental section.

Abstract

This application brief describes the separation of nucleotide phosphates on ACQUITY UPLC BEH Amide column.

Introduction

The nucleotide phosphates used in this study are-

- 1. Adenosine monophosphate (AMP)
- 2. Uridine monophosphate (UMP)
- 3. Adenosine diphosphate (ADP)
- 4. Uridine diphosphate (UDP)
- 5. Adenosine triphosphate (ATP)
- 6. Uridine triphosphate (UTP)




Figure 1. Structures of the compounds used in this study.

Experimental

Test conditions

Column: ACQUITY UPLC BEH Amide, 2.1 x 100 mm, 1.7 μ Part Number: 186004801 Isocratic Mobile Phase: 70/30 ACN/H₂O with 27 mM KH₂PO₄, pH 4.5 Flow Rate: 0.5 mL/min Injection Volume: 5 µL (PLNO) Sample Concentration: shown on chromatogram Sample Diluent: 80/20 ACN/H₂O Column Temperature: 25 °C Weak Needle Wash: 95/5 ACN/H₂O Instrument: Waters ACQUITY UPLC with ACQUITY PDA Detection: UV 260 nm Sampling Rate: 20 Hz

0.1 s

Results and Discussion

Time Constant:




Figure 2. Sample Concentration 1- AMP (50 μ g/mL), 2- UMP (50 μ g/mL), 3- ADP (100 μ g/mL), 4- UDP (100 μ g/mL), 5- ATP (100 μ g/mL), 6- UTP (100 μ g/mL)

Featured Products

ACQUITY UPLC System https://www.waters.com/514207

ACQUITY UPLC PDA Detector https://www.waters.com/514225

WA64069, August 2009

©2019 Waters Corporation. All Rights Reserved.